ELSEVIER

Neuroscience Letters 282 (2000) 203–206

Neuroscience Letters

www.elsevier.com/locate/neulet

Author Index to Volume 282

Abe, K., Saito, H., The mitogen-activated protein kinase cascade mediates neurotrophic effect of epidermal growth factor in cultured rat hippocampal neurons, 89
Akazawa, T, Tokuno, H., Nambu, A., Hamada, I., Ito, Y., Ikeuchi, Y., Imanishi, M., Hasegawa, N., Hatanaka, N., Takada, M., A cortical motor region that represents the cutaneous back muscles in the macaque monkey, 125
Alcayaga, J., Cerpa, V., Retamal, M., Arroyo, J., Iturriaga, R., Zapata, P., Adenosine triphosphate-induced peripheral nerve discharges generated from the cat petrosal ganglion in vitro, 185
Anderson, R.E., see Coert, B.A., 5
Andreu, F., see Zamanillo, D., 169
Antle, M.C., Glass, J.D., Mistlberger, R.E., 5-HT_{1A} autoreceptor antagonist-induced 5-HT release in the hamster suprachiasmatic nuclei: effects on circadian clock resetting, 97
Arakawa, Y., see Watanabe, M., 61
Arroyo, J., see Alcayaga, J., 185
Axelman, K., see Mattila, K.M., 45

Baccari, M.C., Romagnani, P., Calamai, F., Impaired nitrergic relaxations in the gastric fundus of dystrophic (mdx) mice, 105
Bai, S.-H., see Jung, C.-S., 53
Barker, J.L., see Dori, A., 77
Barone, F.C., see Legos, J.J., 189
Becker, J., see Sarco, D.P., 113
Bedford, D.F., Steege, J.W., Lee, W.A., Effects of vision on head stability and torques during voluntary pulls made by standing humans, 9
Beirão, P.S.L., see Cruz, J.S., 25
Blok, B.F.M., Holstege, G., The pontine micturition center in rat receives direct lumbosacral input. An ultrastructural study, 29
Blomberg, M., see Mattila, K.M., 45
Bueno-López, J.L., see Gómez-Urquijo, S.M., 177
Byun, H.-S., see Lee, J.-H., 49

Calamai, F., see Baccari, M.C., 105
Cao, X.-D., see Yu, C.-X., 161
Carlier, P.R., see Liu, J., 165
Cerpa, V., see Alcayaga, J., 185
Chang, Y.-Y., see Liu, J.-S., 85
Chen, S.-S., see Liu, J.-S., 85
Chen, W.-H., see Liu, J.-S., 85
Coert, B.A., Anderson, R.E., Meyer, F.B., Exogenous spermine reduces ischemic damage in a model of focal cerebral ischemia in the rat, 5
Cravatt, B.F., see Egertová, M., 13
Cruz, J.S., Matavel, A.C.S., Leão-Filho, H.M., Moraes-Santos, T., Beirão, P.S.L., Tityustoxin effect on nerve compound action potentials requires extracellular sodium, 25

de Lange, R.P.J., see Ünlü, M., 149
de Silva, R., see Ünlü, M., 149
DeLeo, J.A., Rutkowski, M.D., Gender differences in rat neuropathic pain sensitivity is dependent on strain, 197
Dietrich, P., see Dragatsis, I., 37
Dori, A., Maric, D., Maric, I., Masalha, R., Barker, J.L., Silverman, W.F., Striatal matrix neurons of the rat differentiate in culture from dissociated fetal progenitor cells isolated by buoyant density centrifugation, 77
Dragatsis, I., Dietrich, P., Zeitlin, S., Expression of the Huntingtin-associated protein 1 gene in the developing and adult mouse, 37

Egertová, M., Cravatt, B.F., Elphick, M.R., Fatty acid amide hydrolase expression in rat choroid plexus: possible role in regulation of the sleep-inducing action of oleamide, 13
Eggert, T., see Kharraz-Tavakol, O.D., 33
Elphick, M.R., see Egertová, M., 13
Endo, Y., see Watanabe, M., 61
Erhardt, J.A., see Legos, J.J., 189

Farré, A.J., see Zamanillo, D., 169
Ferriero, D.M., see Sarco, D.P., 113
Fredrikson, M., see Lekander, M., 193
Fujiwara, M., see Taga, G., 101
Fukuda, H., see Nakaso, K., 57
Fukuda, Y., see Morigiwa, K., 153
Fuwa, T., see Miwa, H., 21

Glass, J.D., see Antle, M.C., 97
Gleichmann, M., Weller, M., Schulz, J.B., Insulin-like growth factor-1-mediated protection from neuronal apoptosis is linked to phosphorylation of the pro-apoptotic protein BAD but not to inhibition of cytochrome c translocation in rat cerebellar neurons, 69
Gómez-Urquijo, S.M., Gutiérrez-Ibarluzea, I., Bueno-López, J.L., Reblet, C., Percentage incidence of γ-aminobutyric acid neurons in the claustrum of the rabbit and comparison with the cortex and putamen, 177
Guitart, X., see Zamanillo, D., 169
Guo, Y., see Regan, R.F., 1
Gutiérrez-Ibarluzea, I., see Gómez-Urquijo, S.M., 177

Hamada, I., see Akazawa, T., 125
Han, Y.-F., see Liu, J., 165
Hasegawa, N., see Akazawa, T., 125
Hasegawa, S., see Iwasa, H., 173
Hashimoto, I., Yoshikawa, K., Kimura, T., Somatosensory evoked potentials to a threshold air-puff can predict stimulus detection in human subjects, 181
Hasselmo, M.E., see Linster, C., 157
Hatanaka, N., see Akazawa, T., 125

PII: S0304-3940(00)00945-9

Häussler, C., see Rolke, B., 81
Heil, M., see Rolke, B., 81
Hennighausen, E., see Rolke, B., 81
Ho, A., see Quiñones-Jenab, V., 120
Ho, I.K., see Jang, C.-G., 129
Ho, W.-L., see Liu, J., 165
Holstege, G., see Blok, B.F.M., 29
Honda, S., see Liu, Y., 137
Honda, T., see Song, S., 65
Hong, H.-N., see Lee, J.-H., 49
Horner, R.L., see Liu, D.K-C., 200
Hotta, M., see Kuriyama, H., 109
Huang, C.-Y., see Liu, J.-S., 85

Ikegami, T., see Wang, L., 133
Ikeuchi, Y., see Akazawa, T., 125
Im, J.-O., see Lee, J.-H., 49
Imanishi, M., see Akazawa, T., 125
Inanami, O., see Tsuji, M., 41
Inui, T., Tanaka, S., Okada, T., Nishizawa, S., Katayama, M., Konishi, J., Neural substrates for depth perception of the Necker cube; a functional magnetic resonance imaging study in human subjects, 145
Ishii, T., see Nakaso, K., 57
Ito, Y., see Akazawa, T., 125
Iturriaga, R., see Alcayaga, J., 185
Iwasa, H., Kikuchi, S., Mine, S., Miyagishima, H., Sugita, K., Sato, T., Hasegawa, S., Up-regulation of type II adenylyl cyclase mRNA in kindling model of epilepsy in rats, 173

Jang, C.-G., Ho, I.K., *N*-Methyl-D-aspartate receptor NR1 subunit mRNA level was decreased in rat brain during pentobarbital withdrawal, 129
Jung, C.-S., Lee, S.-J., Paik, S.-S., Bai, S.-H., Run-up of γ-aminobutyric $acid_C$ responses in catfish retinal cone-horizontal cell axon-terminals is modulated by protein kinase A and C, 53

Kalaria, R., see Ünlü, M., 149
Kamitani, T., see Wang, L., 133
Kang, H., Schuman, E.M., Intracellular Ca^{2+} signaling is required for neurotrophin-induced potentiation in the adult rat hippocampus, 141
Kao, Y.-F., see Liu, J.-S., 85
Katayama, M., see Inui, T., 145
Katoh-Semba, R., see Watanabe, M., 61
Kawato, S., see Mukai, H., 93
Kharraz-Tavakol, O.D., Eggert, T., Mai, N., Straube, A., Learning to write letters: transfer in automated movements indicates modularity of motor programs in human subjects, 33
Kikuchi, S., see Iwasa, H., 173
Kim, D., see Lee, J.-H., 49
Kimoto, K., see Watanabe, M., 61
Kimura, K., see Nakaso, K., 57
Kimura, T., see Hashimoto, I., 181
Kitayama, M., see Nakaso, K., 57
Kohsaka, S., see Liu, Y., 137
Koizumi, H., see Taga, G., 101
Konishi, J., see Inui, T., 145
Konishi, Y., see Taga, G., 101
Kreek, M.J., see Quiñones-Jenab, V., 120
Krey, L.C., see Quiñones-Jenab, V., 120
Kumar, N., see Regan, R.F., 1
Kuriyama, H., Hotta, M., Wakabayashi, I., Shibasaki, T., A 6-day intracerebroventricular infusion of the growth hormone-releasing peptide KP-102 stimulates food intake in both non-stressed and intermittently-stressed rats, 109
Kuroiwa, Y., see Wang, L., 133
Kuwabara, M., see Tsuji, M., 41

Laippala, P., see Mattila, K.M., 45
Lan, M.-Y., see Liu, J.-S., 85
Lannfelt, L., see Mattila, K.M., 45
Leão-Filho, H.M., see Cruz, J.S., 25
Lee, J.-H., Hong, H.-N., Im, J.-O., Byun, H.-S., Kim, D., The formation of PHF-1 and SMI-31 positive dystrophic neurites in rat hippocampus following acute injection of okadaic acid, 49
Lee, N.T.-K., see Liu, J., 165
Lee, S.-J., see Jung, C.-S., 53
Lee, W.A., see Bedford, D.F., 9
Legos, J.J., Whitmore, R.G., Erhardt, J.A., Parsons, A.A., Tuma, R.F., Barone, F.C., Quantitative changes in interleukin proteins following focal stroke in the rat, 189
Lehtimäki, T., see Mattila, K.M., 45
Lekander, M., Fredrikson, M., Wik, G., Neuroimmune relations in patients with fibromyalgia: a positron emission tomography study, 193
Li, M., see Wang, L., 133
Li, Z.-q., Yamamoto, Y., Morimoto, T., Ono, J., Okada, S., Yamatodani, A., The effect of pentylenetetrazole-kindling on the extracellular glutamate and taurine levels in the frontal cortex of rats, 117
Linster, C., Hasselmo, M.E., Neural activity in the horizontal limb of the diagonal band of Broca can be modulated by electrical stimulation of the olfactory bulb and cortex in rats, 157
Liu, D.K-C., Horner, R.L., Wojtowicz, J.M., Time of day determines modulation of synaptic transmission by adenosine in the rat hippocampal slices, 200
Liu, J., Ho, W.-L., Lee, N.T.-K., Carlier, P.R., Pang, Y.-P., Han, Y.-F., *Bis*(7)-tacrine, a novel acetylcholinesterase inhibitor, reverses AF64A-induced deficits in navigational memory in rats, 165
Liu, J.-S., Chang, Y.-Y., Wu, H.-S., Huang, C.-Y., Chen, W.-H., Lan, M.-Y., Kao, Y.-F., Chen, S.-S., Transtentorial cerebellar c-*jun* expression after focal cerebral cortical injury in mice
Liu, Y., Honda, S., Kohsaka, S., Nakajima, K., Plasminogen enhances the secretion of plasminogen activator inhibitor-1 from cultured rat astrocytes, 137

Mai, N., see Kharraz-Tavakol, O.D., 33
Maki, A., see Taga, G., 101
Maric, D., see Dori, A., 77
Maric, I., see Dori, A., 77
Masalha, R., see Dori, A., 77
Matavel, A.C.S., see Cruz, J.S., 25
Matsubara, S., see Wang, L., 133
Mattila, K.M., Axelman, K., Rinne, J.O., Blomberg, M., Lehtimäki, T., Laippala, P., Röyttä, M., Viitanen, M., Wahlund, L.-O., Winblad, B., Lannfelt, L., Interaction between estrogen receptor 1 and the ε4 allele of apolipoprotein E increases the risk of familial Alzheimer's disease in women, 45
Meyer, F.B., see Coert, B.A., 5
Mine, S., see Iwasa, H., 173
Mistlberger, R.E., see Antle, M.C., 97
Miwa, H., Nishi, K., Fuwa, T., Mizuno, Y., Effects of blockade of metabotropic glutamate receptors in the subthalamic nucleus on haloperidol-induced Parkinsonism in rats, 21

Miyagishima, H., see Iwasa, H., 173
Mizuno, Y., see Miwa, H., 21
Moraes-Santos, T., see Cruz, J.S., 25
Morigiwa, K., Quan, M.-z., Murakami, M., Yamashita, M., Fukuda, Y., P2 purinoceptor expression and functional changes of hypoxia-activated cultured rat retinal microglia, 153
Morimoto, T., see Li, Z.-q., 117
Mukai, H., Uchino, S., Kawato, S., Effects of neurosteroids on Ca^{2+} signaling mediated by recombinant *N*-methyl-D-aspartate receptor expressed in Chinese hamster ovary cells, 93
Murakami, M., see Morigiwa, K., 153

Nakajima, K., see Liu, Y., 137
Nakashima, K., see Nakaso, K., 57
Nakaso, K., Kitayama, M., Fukuda, H., Kimura, K., Yanagawa, T., Ishii, T., Nakashima, K., Yamada, K., Oxidative stress-related proteins A170 and heme oxygenase-1 are differently induced in the rat cerebellum under kainate-mediated excitotoxicity, 57
Nambu, A., see Akazawa, T., 125
Netzer, R., see Rundfeldt, C., 73
Nishi, K., see Miwa, H., 21
Nishizawa, S., see Inui, T., 145

Ohba, M., see Song, S., 65
Okada, S., see Li, Z.-q., 117
Okada, T., see Inui, T., 145
Ono, J., see Li, Z.-q., 117
Ovalle, S., see Zamanillo, D., 169

Packard, M.G., see Schroeder, J.P., 17
Paik, S.-S., see Jung, C.-S., 53
Palmer, C., see Sarco, D.P., 113
Pang, Y.-P., see Liu, J., 165
Parsons, A.A., see Legos, J.J., 189
Pérez, M.P., see Zamanillo, D., 169

Quan, M.-z., see Morigiwa, K., 153
Quiñones-Jenab, V., Krey, L.C., Schlussman, S.D., Ho, A., Kreek, M.J., Chronic 'binge' pattern cocaine alters the neuroendocrine profile of pregnant rats, 120

Reblet, C., see Gómez-Urquijo, S.M., 177
Regan, R.F., Guo, Y., Kumar, N., Heme oxygenase-1 induction protects murine cortical astrocytes from hemoglobin toxicity, 1
Retamal, M., see Alcayaga, J., 185
Rinne, J.O., see Mattila, K.M., 45
Rolke, B., Heil, M., Hennighausen, E., Häussler, C., Rösler, F, Topography of brain electrical activity dissociates the sequential order transformation of verbal versus spatial information in humans, 81
Romagnani, P., see Baccari, M.C., 105
Romero, G., see Zamanillo, D., 169
Rösler, F, see Rolke, B., 81
Röyttä, M., see Mattila, K.M., 45
Rundfeldt, C., Netzer, R., The novel anticonvulsant retigabine activates M-currents in Chinese hamster ovary-cells tranfected with human KCNQ2/3 subunits, 73
Rutkowski, M.D., see DeLeo, J.A., 197

Saito, H., see Abe, K., 89
Saito, Y., see Song, S., 65
Sarco, D.P., Becker, J., Palmer, C., Sheldon, R.A., Ferriero, D.M., The neuroprotective effect of deferoxamine in the hypoxic-ischemic immature mouse brain, 113
Sato, T., see Iwasa, H., 173
Schlussman, S.D., see Quiñones-Jenab, V., 120
Schroeder, J.P., Packard, M.G., Role of dopamine receptor subtypes in the acquisition of a testosterone conditioned place preference in rats, 17
Schulz, J.B., see Gleichmann, M., 69
Schuman, E.M., see Kang, H., 141
Sheldon, R.A., see Sarco, D.P., 113
Shibasaki, T., see Kuriyama, H., 109
Silverman, W.F., see Dori, A., 77
Song, S., Ohba, M., Saito, Y., Honda, T., Takashima, A., Takahashi, H., Proteolytic processing and degradation of human presenilin-1 expressed in yeast, 65
St. Clair, D., see Ünlü, M., 149
Steege, J.W., see Bedford, D.F., 9
Straube, A., see Kharraz-Tavakol, O.D., 33
Sugita, K., see Iwasa, H., 173
Suzuki, Y., see Wang, L., 133

Tachibana, T., see Taga, G., 101
Taga, G., Konishi, Y., Maki, A., Tachibana, T., Fujiwara, M., Koizumi, H., Spontaneous oscillation of oxy- and deoxy- hemoglobin changes with a phase difference throughout the occipital cortex of newborn infants observed using non-invasive optical topography, 101
Takada, M., see Akazawa, T., 125
Takahashi, H., see Song, S., 65
Takahashi, T., see Wang, L., 133
Takashima, A., see Song, S., 65
Tanaka, S., see Inui, T., 145
Tokuno, H., see Akazawa, T., 125
Tsuji, M., Inanami, O., Kuwabara, M., Neuroprotective effect of α-phenyl-N-*tert*-butylnitrone in gerbil hippocampus is mediated by the mitogen-activated protein kinase pathway and heat shock proteins, 41
Tuma, R.F., see Legos, J.J., 189

Uchino, S., see Mukai, H., 93
Ünlü, M., de Lange, R.P.J., de Silva, R., Kalaria, R., St. Clair, D., Detection of complement factor B in the cerebrospinal fluid of patients with cerebral autosomal dominant arteriopathy with subcortical infarcts and leukoencephalopathy disease using two-dimensional gel electrophoresis and mass spectrometry, 149

Viitanen, M., see Mattila, K.M., 45

Wahlund, L.-O., see Mattila, K.M., 45
Wakabayashi, I., see Kuriyama, H., 109
Wang, J., see Wang, L., 133
Wang, L., Kuroiwa, Y., Li, M., Kamitani, T., Wang, J., Takahashi, T., Suzuki, Y., Ikegami, T., Matsubara, S., The correlation between P300 alterations and regional cerebral blood flow in non-demented Parkinson's disease, 133
Watanabe, M., Endo, Y., Kimoto, K., Katoh-Semba, R., Arakawa, Y., Inhibition of adjuvant-induced inflammatory hyperalgesia in rats by local injection of neurotrophin-3, 61
Weller, M., see Gleichmann, M., 69
Whitmore, R.G., see Legos, J.J., 189
Wik, G., see Lekander, M., 193

Winblad, B., see Mattila, K.M., 45
Wojtowicz, J.M., see Liu, D.K-C., 200
Wu, G.-C., see Yu, C.-X., 161
Wu, H.-S., see Liu, J.-S., 85

Xu, S.-F., see Yu, C.-X., 161

Yamada, K., see Nakaso, K., 57
Yamamoto, Y., see Li, Z.-q., 117
Yamashita, M., see Morigiwa, K., 153
Yamatodani, A., see Li, Z.-q., 117
Yanagawa, T., see Nakaso, K., 57
Yoshikawa, K., see Hashimoto, I., 181
Yu, C.-X., Zhu, C.-B., Xu, S.-F., Cao, X.-D., Wu, G.-C., Selective MT_2 melatonin receptor antagonist blocks melatonin-induced antinociception in rats, 161

Zamanillo, D., Andreu, F., Ovalle, S., Pérez, M.P., Romero, G., Farré, A.J., Guitart, X., Up-regulation of $sigma_1$ receptor mRNA in rat brain by a putative atypical antipsychotic and sigma receptor ligand, 169
Zapata, P., see Alcayaga, J., 185
Zeitlin, S., see Dragatsis, I., 37
Zhu, C.-B., see Yu, C.-X., 161